

Frequency of Celiac Disease in Patients Presenting with Nutritional Anemia

Ammar Asghar, Muhammad Hassan Zafar, Salman Munir

ABSTRACT

Introduction: Celiac disease (CD) is a gluten sensitive enteropathy characterized by villous atrophy of the small intestine in genetically susceptible individuals. Anemia is frequently seen in celiac disease owing to greatly reduced iron, folic acid and B12 absorption. **Objectives:** To determine the frequency of celiac disease in patients presenting with nutritional anemias. **Study design:** Descriptive cross-sectional study **Setting:** department of Internal Medicine, Services Hospital Lahore. **Duration of study:** Seven months from May 2017 to December 2017. **Sample size:** The calculated sample size is 100 cases. **Sampling technique:** Non-probability purposive sampling. **Data collection procedure:** 100 consecutive patients presenting with nutritional anemia were enrolled. Complete hematologic workup was done including blood complete picture (CP), peripheral smear, iron, folate and B12 levels. Anemia was defined as Hb <13 g/dL in males and <12 g/dL in females. Only cases with anemia secondary to iron, folate or B12 deficiency i.e. nutritional causes were included in the study. Patients were then subjected to anti-tTG antibody testing. Jejunal biopsy was taken in all those cases which tested positive for anti tTG antibody. A diagnosis of CD was established upon positive histopathology findings. All the data was recorded on a pre-designed proforma and analyzed by SPSS version 21.0. **Results:** The prevalence of Celiac disease was found to be 12%. 10 out of 12 celiac disease cases were having IDA; 1 each showed dimorphic picture and concomitant B12 + folate deficiency respectively with none of the CD patients having folate or B12 deficiency alone. **Conclusion:** Celiac disease may be found in up to 12% of Pakistanis presenting with nutritional anemia. Hence, all those patients not responding to standard anemia treatment should be screened for celiac disease.

Keywords: Anti-tissue transglutaminase antibody (Anti-tTG), Celiac disease (CD), Iron deficiency anemia (IDA)

Corresponding Author

Dr. Ammar Asghar

House officer, Medicine
Services Hospital, Lahore
Contact: +92 300-9676452
Email: umairasghar51@yahoo.com

Submitted for Publication: 13-04-2018

Accepted for Publication: 27-05-2018

Article Citation: Asghar A, Zafar MH, Munir S. Frequency of Celiac Disease in Patients Presenting with Nutritional Anemia. APMC 2018;12(2):151-3.

INTRODUCTION

Celiac disease (CD) is a gluten sensitive enteropathy characterized by villous atrophy of the small intestine in genetically susceptible individuals. Its worldwide prevalence is estimated to be somewhere between 0.5-1%.¹ The exact prevalence of celiac disease in Pakistan is unknown. HLA DQ2 and DQ8 subtypes genetically predispose the individual to celiac disease.²

The pathophysiology behind celiac disease involves abnormal interaction between gluten, immune system and the gut leading to destruction of villi. Villous atrophy in celiac disease renders absorption of various nutrients ineffective. Iron, B12, folic acid and fat-soluble vitamin deficiencies are particularly noticeable.³ Patients with classical celiac disease present with gastrointestinal symptoms of malabsorption i.e. diarrhea, abdominal pain and weight loss. However, majority of the cases are asymptomatic. Estimates show that up to 90% of the cases remain undiagnosed and the classic cases just represent the tip of iceberg.⁴ Diagnosis of celiac disease involves antibody testing and histopathology. Anti-Tissue-transglutaminase antibody is the best initial test. A positive test has to be confirmed with jejunal biopsy which is the gold standard investigation for diagnosis of celiac disease.⁵

Anemia is frequently seen in celiac disease owing to greatly reduced iron, folic acid and B12 absorption. It has been estimated that iron deficiency anemia is present in up to 80 to 90% of patients with celiac disease.⁶ What is important to recognize is the fact that sometimes the sole presentation of celiac disease is that of anemia without any diarrheal symptoms.⁷ This IDA is unique in a way that it does not respond to iron supplementation therapy as the absorptive surface areas has been greatly reduced. Its management involves gluten free diet and can take up to 1 year for complete recovery. Presence of a dimorphic picture on blood smear is highly suggestive of concomitant iron and folate / B12 deficiency.⁸ As vitamin B12 is absorbed in the terminal ileum, it is hard to explain its deficiency in celiac disease. Dahele et al⁹ reported that 56% of the anemic patients 30% of the nonanemic patients with CD were vitamin B12-deficient.

A recent study by Shahzad et al¹⁰ reported a prevalence of 12.99% in IDA patients. An Iranian study by Baghbanian et al¹¹ reported a prevalence of 10.4% amongst IDA patients. Kavimandan et al⁶ found out CD was prevalent 10.42% of the patients with nutritional anemia. Local studies are lacking in this area so we decided to conduct this study with the principal aim of determining the prevalence of CD in patients with nutritional anemia.

METHODOLOGY

Study design: Descriptive cross-sectional study.

Setting: department of Internal Medicine, services hospital Lahore.

Duration of study: Seven months from May 2017 to December 2017.

Sample size: The sample size was calculated using Open Epi calculator with the statistical assumptions of 6% alpha error and 95 % confidence interval taking prevalence of celiac disease to be 10.42%⁶ amongst patients with nutritional anemia and comes out to be at least 100 patients for this study.

Sampling technique: Non probability purposive sampling. After getting ethical approval from IRB, 100 consecutive patients presenting with nutritional anemia were enrolled. Patients having malignancy, chronic diseases (IBD, tuberculosis, autoimmune diseases), pregnancy or any primary hematological disorder were excluded from the study. Informed consent was taken in each case. Anemia was defined as Hb <13 g/dL in males and <12 g/dL in females. Only cases with anemia secondary to iron, folate or B12 deficiency i.e. nutritional causes were included in the study. Detailed clinical history was taken from each patient. Complete hematologic workup was done including blood CP, peripheral smear, iron, folate and B12 levels. Patients were then subjected to certain screening and diagnostic tests. Blood sample was withdrawn from a peripheral vein followed by testing for Anti-tissue transglutaminase antibodies (tTG) antibody. Jejunal biopsy was taken in all those cases which tested positive for anti tTG antibody. The specimens were inspected by a histopathologist having experience of at least 5 years. A diagnosis of CD was established upon positive histopathology findings of crypt hyperplasia and villous atrophy.

All the data was recorded on a pre-designed proforma and analyzed by SPSS version 21.0. Mean and standard deviation was calculated for all quantitative variables like age and Hb level etc. Frequency and percentage was calculated for all qualitative variables like gender, cause of nutritional anemia, prevalence of celiac disease etc. Data was represented as pie charts and bar graphs.

RESULTS

100 patients took part in the study comprising of 45 males and 55 females. Mean age of the patients was 34.12 ± 11.23 years. A mean Hb of 8.71 ± 1.23 g/dL was observed (Table 1). The prevalence of Celiac disease was found to be 12% (Figure 1). Initially 14 patients tested positive for anti-tTG antibody. Upon further testing only 12 of them showed the characteristic biopsy findings necessary to qualify for diagnosis of celiac disease. IDA was the most frequent cause of anemia present in 85% of the cases. 5 patients had folate deficiency; 3 had B12 deficiency whereas 6 patients showed concomitant B12 and folate deficiency. One patient showed dimorphic blood picture. 10 out of 12 celiac disease cases were having IDA; 1 each showed dimorphic picture and concomitant B12 + folate deficiency

respectively with none of the CD patients having folate or B12 deficiency alone (Figure 2).




Figure 1: Frequency of celiac disease in patients with nutritional anemia




Figure 2: Distribution of celiac disease among various nutritional anemias

Table 1: Baseline Characteristics

Patients Characteristics	N=100 (%)
Mean age \pm SD	34.12 \pm 11.23 years
Mean Hb \pm SD	8.71 \pm 1.03 g/dL
Males / Females	45 / 55
Positive Serology	14 (14%)
Positive Jejunal biopsy	12 (12%)
IDA / Folate deficiency anemia / B12 deficiency anemia / Concomitant B12 + Folate deficiency / Dimorphic anemia	85 / 5 / 3 / 6 / 1

DISCUSSION

Anemia is one of the common extraintestinal manifestation of celiac disease. Often, the only symptoms present are those of anemia without any characteristic features of celiac disease like diarrhea etc. This anemia is resistant to standard treatment because the problem lies in the absorptive surface leading to reduced absorption of vital nutrients. The only possible cure is gluten free diet. Therefore, it is important to screen all those patients with anemia who do not seem to respond to routine management. Hence we conducted this study to determine the prevalence of CD in anemia patients in our local population.

Our study reported a prevalence of 12% for CD amongst nutritional anemia patients. This was comparable to the findings of Kavimandan et al⁶ who reported a prevalence of 10.42%. Some other studies conducted on IDA patients reported an overall prevalence of 12.99% and 10.4% respectively.⁹⁻¹⁰ Our study showed that 10 out of 85 IDA patients (11.76%) were having celiac disease. Mean age of patients in our study was 34.12±11.23 years which was quite close to that reported by Kavimandan et al⁶ i.e. 32.1±13.1 years. In our study, 85% of the subjects had IDA. This was consistent with the findings of Kavimandan et al⁶ who showed that 83.3% of patients were having IDA. Megaloblastic anemia was seen in 14% (5%: Folate; 3%: B12; 6%: concomitant B12+Folate deficiency) patients which was comparable to the findings of Kavimandan et al who reported a frequency of 11.46%. Mean Hb observed in our study was 8.71±1.03 g/dL. This was quite similar to the value reported by Shahzad et al⁹ who reported a mean Hb of 8.81±1.23 g/dL.

We relied upon jejunal biopsy findings in order to diagnose the patient with celiac disease. However, we did not take biopsy of all the 100 patients as it was not feasible. Instead we screened the patients using serologic testing for anti-tTG antibody and only performed jejunal biopsy on patients who tested positive on serology. 14 of the patients screened positive for celiac disease initially however the diagnosis could be validated in only 12 of those 14 cases. A similar trend was noticed by Kavimandan et al.⁶

There were certain limitations to our study. Effect modifiers and confounders (gender, cause of nutritional anemia) were not controlled through stratification. The sample size was small and future large-scale surveys are warranted to estimate the true prevalence of CD amongst anemia patients and screening guidelines updated accordingly.



CONCLUSION

Celiac disease may be found in up to 12% of Pakistanis presenting with nutritional anemia. Hence, all those patients not responding to standard anemia treatment should be screened for celiac disease.

REFERENCES

1. Gujral N, Freeman HJ, Thomson AB. Celiac disease: prevalence, diagnosis, pathogenesis and treatment. *World J Gastroenterol.* 2012;18(42):6036.
2. Megiorni F, Pizzuti A. HLA-DQA1 and HLA-DQB1 in Celiac disease predisposition: practical implications of the HLA molecular typing. *J Biomed Sci.* 2012;19:88.
3. Garcia-Manzanares Á, Lucendo AJ. Nutritional and dietary aspects of celiac disease. *Nutr Clin Pract.* 2011;26(2):163-73.
4. Rashid M, Khan AG. Celiac disease in Pakistan: challenges and opportunities. *J Ayub Med Coll Abbottabad.* 2009;21(3):1-2.
5. Guandalini S, Assiri A. Celiac disease: a review. *JAMA Pediatr.* 2014;168(3):272-8.
6. Kavimandan A, Sharma M, Verma AK, Das P, Mishra P, Sinha S, Mohan A, Sreenivas V, Gupta SD, Makharia GK. Prevalence of celiac disease in nutritional anemia at a tertiary care center. *Indian J Gastroenterol.* 2014;33(2):114-8.
7. Brandimarte G, Tursi A, Giorgetti GM. Changing trends in clinical form of celiac disease. Which is now the main form of celiac disease in clinical practice?. *Minerva Gastroenterol Dietol.* 2002;48(2):121-30.
8. Halfdanarson TR, Litzow MR, Murray JA. Hematologic manifestations of celiac disease. *Blood.* 2007;109(2):412-21.
9. Dahele A, Ghosh S. Vitamin B12 deficiency in untreated celiac disease. *Am J Gastroenterol* 2001;745-50.
10. Shahzad A, Sahto AA, Samina. Frequency of celiac disease; patients presenting with iron deficiency anemia at tertiary care hospital. *Professional Med J.* 2016;23(7):812-6.
11. Baghbanian M, Farahat A, Vahedian Ha, Sheyda E, Zare-khormizi MR. The prevalence of celiac disease in patients with iron-deficiency anemia in center and south area of Iran. *Arq Gastroenterol.* 2015;52(4):278-82.

AUTHORSHIP AND CONTRIBUTION DECLARATION

AUTHORS	Contribution to The Paper	Signatures
Dr. Ammar Asghar House officer, Medicine Services Hospital, Lahore	Writing the Manuscript	
Dr. Muhammad Hassan Zafar House Officer, Medicine Services Hospital, Lahore	Data Collection, References writing	
Dr. Salman Munir Assistant Professor, Medicine Punjab Institute of Cardiology, Lahore	Proof Reading, Statistical Analysis	